

PUBLIC HEALTH STATISTICS

STATE OF

OKLAHOMA

1954



PART 1

REPORTABLE DISEASES

PUBLIC HEALTH STATISTICS

STATE OF

OKLAHOMA

1954



PART I

REPORTABLE DISEASES

Oklahoma State Department of Health
Oklahoma City, Oklahoma
G. F. MATHEWS, M. D., Commissioner

TABLE OF CONTENTS

	Page
Discussion.....	1
Diphtheria.....	2
Intestinal Diseases.....	2
Infectious Hepatitis.....	3
Malaria.....	3
Measles.....	5
Meningococcal Infections.....	5
Polioomyelitis.....	5
Respiratory Streptococcal Infections.....	6
Whooping Cough.....	7
Other Acute Communicable Diseases.....	7
Rheumatic Fever.....	7
Tuberculosis.....	8
Veneral Diseases.....	10
Malignant Neoplasms.....	11
Appendix	
Table I. Reported Cases of Selected Communicable Diseases, Number and Rate (Number per 100,000 Estimated Population), Oklahoma, 1945-1954.....	15
Table II. Reported Cases of Communicable Diseases, Number and Rate (Number per 100,000 Estimated Population), by Race, Oklahoma, 1954.....	16
Table III. Reported Cases of Communicable Diseases, by Month, Oklahoma, 1954.....	16
Table IV. Reported Civilian Cases of Communicable Diseases, Number and Rate (Number per 100,000 Estimated Population), and Number by Race, by Urban and Rural Residence, Oklahoma, 1954.....	17
Table V. Reported Cases of Malignant Neoplasms, by Primary Site of Lesion, Race and Sex, Oklahoma, 1954.....	17
Table VI. Reported Cases of Selected Communicable Diseases by Sex and Race, Oklahoma, 1954.....	18
Table VII. Reported Cases of Selected Communicable Diseases by Age, Oklahoma, 1954.....	19
Table VIII. Reported Cases of Communicable Diseases by County of Residence, Oklahoma City, Tulsa City, and Military, 1954.....	20

PUBLIC HEALTH STATISTICS OF OKLAHOMA
REPORTABLE DISEASES

1954

Statistical data pertaining to the cases of diseases reported to the State Department of Health during the year 1954 have been tabulated for this eleventh edition of Public Health Statistics of Oklahoma, Part I. Reports of cases on prescribed forms are received weekly from doctors, local health departments, hospitals, and clinics. In addition, cases of some of the more serious diseases in persons who supposedly contracted the disease in Oklahoma but are under medical care in other states are received through interstate reciprocal notifications. Laboratory reports from the State Laboratory which are positive for certain organisms are queried for additional information but are not counted as reported cases unless the diagnoses are confirmed by the attending physician. Cases of selected diseases reported by death certificates are included as reported cases also, provided, of course, they were not previously reported through regular channels.

In so far as possible, duplicate reports are eliminated; files for the acute communicable diseases are maintained for the year and cumulative files are kept for the long-term or recurrent diseases. Allocation of cases among the civilian population to counties and cities is determined by the place where the disease was presumably contracted, which in most instances is the same as the place of residence. Military cases, however, although included in State totals and tabulated separately, are not allocated to any county or city since the control of communicable diseases among the military population is the responsibility of the military officials and not of the local health authorities.

Eight master tables in the appendix show pertinent information regarding the race, sex, age, month of report, and county of occurrence of the reported cases for each disease. Observations of particular interest have been selected for discussion, and tables and graphs have been used for illustration in the narrative. Population estimates made by the Division of Statistics have been used for computing rates; all incidence rates represent the number of cases per 100,000 estimated population within the specific classification. The number of deaths for the year used for computing case fatality rates for certain diseases are provisional figures for 1954; final mortality data will be published in Part II of Public Health Statistics of Oklahoma.

Although it is a recognized fact that the reporting of disease is not complete, it is difficult to determine the extent of underreporting. It is assumed, however, that the completeness of reporting would vary from disease to disease according to the severity of the disease, public interest in the disease, and immediate need for the institution of treatment or control measures for the disease. The number of cases of a particular disease reported by death certificate only is one readily available measure of the extent of underreporting. Some of these numbers and their ratio to the total case reports for the selected disease are shown in Table 1.

Table 1
Cases of Diseases Reported by Death Certificate Only
Oklahoma, 1954

Disease	Total Number Reported Cases	Cases Reported by Death Certificate	Per Cent Reported by Death Certificate
Diphtheria	42	-	-
Dysentery	207	10	4.8
Encephalitis, infectious	24	4	16.7
Hepatitis, infectious	347	8	2.3
Meningococcal infections	58	2	3.4
Poliovellitis, acute	559	2	0.4
Rheumatic fever	52	9	17.3
Rocky Mountain spotted fever	6	-	-
Septic sore throat	566	4	0.7
Tetanus	7	3	42.9
Tuberculosis, all forms	1,642	65	4.0
Whooping cough	219	3	1.4

DIPHTHERIA

The number of cases of diphtheria continued to decrease, with only 42, or 1.9 cases per 100,000 population, reported during the year 1954. Three cases ended in death, however, following a year in which no deaths from the disease had been recorded.

Of the total cases, 20 or 47.6 per cent, were under ten years of age. Thirty-six of the cases were white, 5 Negro and one Indian.

INTESTINAL DISEASES

During the year, 207 cases of dysentery were reported, only a few less than were reported in the previous year when 225 cases were recorded. Most of the cases of dysentery occurred in children, with 20.9 per cent of the cases with age specified reported as under one year of age and 41.8 ages 1 through 9; only 37.2 per cent were 10 years of age or older.

Over 75 per cent of the dysentery cases were specified as bacillary. The attack rate was very high among Indians, 17.1 per 100,000 population, as compared to the rates for the white and Negro populations, 2.7 and 3.4, respectively. Twenty-four of the dysentery cases were specified as amebic and 27 were other forms or unspecified as to type.

An increase in the number of typhoid fever cases, 63, reported in 1954 over the previous year's all-time low record of 46, was recorded. The rate of 2.8 cases per 100,000 estimated population, however, is the same as the previous five-year average rate. The rate was high for the Indian population, 20.5, and low for the Negro, 0.7, compared to the rate of 2.5 for the white population. In the one month of August, 19 cases were reported, bringing the total for the summer (June, July, and August) to 30, or almost half of the cases for the year. Three deaths during 1954 were attributed to typhoid fever, making a case fatality rate of 4.8 per cent. Only two new typhoid carriers were reported during the year, bringing the total of known carriers up to 37.

During the year, 55 cases of food poisoning were reported as compared to 36 in the previous year. Thirty-five of the cases were specified as salmonella infections, 13 as staphylococcal poisoning, and the other 7 were unclassified. Only one outbreak of any size was reported when 24 cases of salmonella infection developed following a civic organization dinner. No cases of botulism were reported.

Brucellosis, another disease commonly arising in the intestinal tract, was reported in 40 cases during 1954, only 2 more than in 1953. The rate of 1.8 cases per 100,000 population, however, was lower than the average rate of 3.7 for the previous five-year period.

The number of paratyphoid fever cases dropped to 38 from the very high number of 68 reported in 1953. The attack rate for paratyphoid fever shows no definite pattern of change. The rate was 1.9 in 1947, 1.7 in 1952, 3.0 in 1953, and 1.7 in 1954; for all other years during the last ten, the rates have not exceeded 0.6.

Only 4 diarrhea of the newborn cases were reported during 1954.

INFECTIOUS HEPATITIS

The number of reported cases of infectious hepatitis, 347, for the year 1954, the fourth year for which figures are available in Oklahoma, almost doubled the number reported in the previous year when 175 were recorded. The rate was highest for the Indian population, 29.8, and lowest for the Negro, 6.2, as compared to the average and white rate of 15.5 per 100,000 population. The peak month was March, with 63 cases reported during that month; 61 per cent of the cases were reported in the period January-May. The case fatality rate, based on 9 deaths and the 347 cases, was 2.6 per cent.


Over two-thirds of the cases, 70.1 per cent, of those with age specified, were under 20 years of age. Chart 1 below shows the attack rate for each age group based on the average number of reported cases for the last four years.

Chart 1
Attack Rates from Infectious Hepatitis, by Age
Oklahoma, 1951-1954



Infectious hepatitis is so new to the list of reportable diseases that a definite trend line has not yet been set in Oklahoma. Although the attack rate appears to be increasing rapidly, part of this increase at least may be due to improvement in reporting during the last two years. The rates for Oklahoma are compared with those for the United States as a whole on Chart 2.

Chart 2
Attack Rates from Infectious Hepatitis
United States and Oklahoma
1947-1954



MALARIA

Only 19 cases of malaria acquired in the United States were reported during the year 1954. Seven other cases were reported as having been acquired outside the United States. Fifteen of the total 26 cases were reported during the month of July. Tulsa County reported the largest incidence, 13 cases acquired in the United States and 3 outside the United States.

MEASLES

The incidence of measles during 1954 did not reach epidemic proportions but was reported in moderate numbers with a total of 3,403 cases recorded, making a rate of 152.1 per 100,000 population. The last peak year was in 1951 when 8,000 cases were reported, giving an attack rate of 358.0. Only 4 deaths during the year 1954 were attributed to measles.

MENINGOCOCCAL INFECTIONS

The number of reported cases of meningococcal infections was 58, making an attack rate of 2.6. The highest rate ever recorded in Oklahoma was 9.5 in 1936, when 223 cases were reported. The lowest rate was recorded in 1941, when only 29 cases were reported, giving an attack rate of 1.2.

Most of the 1954 cases were in children, with 49, or 84.5 per cent, of the cases under 15 years of age. Ten deaths during the year were attributed to meningococcal infections, 17.2 per cent of the reported cases. Although the attack and death rates from meningococcal infections are low, the case fatality rate from this disease is extremely high in comparison to some of the other case fatality rates for diseases most frequently reported in children and adolescents. Three-year average case fatality rates for some of these diseases are shown in Table 2.

Table 2
Case Fatality Rates for Selected Communicable Diseases
Oklahoma, 1952-1954

Disease	Reported Cases	Deaths	Case Fatality Rate
Scarlet fever and streptococcal sore throat	2,907	14	0.5
Diphtheria	185	5	2.7
Meningococcal infections	168	39	23.2
Polioviruses	2,206	84	3.8
Measles	8,920	23	0.3
Infectious hepatitis	592	28	4.7
Whooping cough	847	22	2.6

POLIOMYELITIS

A total of 559 cases of acute poliomyelitis was reported during 1954. The attack rate for the population as a whole was 25.0, for the white population, 26.2, for the Indian, 20.5, and for the Negro, 6.9.

Over half of the case reports did not specify whether or not the case was paralytic or non-paralytic. Of those that did, 52.1 per cent were reported as paralytic. Table 3 below shows the age distribution of the cases for which the paralytic status was known. The proportion of the reports that included the paralytic status had increased slightly over the 1953 reporting. In 1952, this information was reported only on 6.5 per cent of the cases, but in 1953 the data were specified for 42.2 per cent of the cases and in 1954 for 42.9 per cent.

Table 3

Reported Cases of Poliomyelitis Specified as Paralytic and Non-Paralytic, by Age, Oklahoma, 1954

Age	Poliomyelitis Cases		Age	Poliomyelitis Cases	
	Paralytic	Non-Paralytic		Paralytic	Non-Paralytic
All ages	125	115	6	7	8
Under 1	5	3	7	6	1
1	5	4	8	4	1
2	9	8	9	17	18
3	7	3	10-14	10	12
4	3	7	15-19	11	15
5	9	7	20-24	26	17
			25 and Over		

July and August were the months of highest incidence, with 159 and 134 cases, respectively, reported during those two months - approximately 52.4 per cent of the cases for the year. The case fatality rate for the year was 4.1 per cent, based on 23 deaths and the 559 reported cases.

RESPIRATORY STREPTOCOCCAL INFECTIONS

The number of cases of both scarlet fever and septic sore throat, 736 and 566, respectively, increased over the 1953 figures, 684 and 262. The combined attack rate for the two was 58.2 cases per 100,000 population, 57.7 for the white population, 21.3 for the Negro, and 27.9 for the Indian. About 77.5 per cent of the cases with age specified were under fifteen years of age. Six deaths during the year were assigned to respiratory streptococcal infections.

WHOOPING COUGH

Whooping cough cases totaled 219 for the year, making an attack rate of 9.8, the lowest on record since 1939 when the rate was 9.1 for 214 cases. Rates among the Negro and Indian populations, 18.6 and 20.5, respectively, were higher than the white rate of 8.5. Of the 40 cases that occurred in infants under one year of age, 6 resulted in death; no other whooping cough deaths occurred during the year. About 67.1 per cent of the cases with age specified were under ten years of age.

OTHER ACUTE COMMUNICABLE DISEASES

No cases of smallpox have been reported since 1950 when 5 cases were recorded.

During the year 1954, no cases of rabies in man were reported either. A total of 54 positive reports for rabies in animals was received for the year. All but 3 of the reports were from the State Laboratory and of those, 49 were dog heads and 2 were cat heads.

Infectious encephalitis was reported in 24 cases and 10 deaths were assigned to the disease. Trichinosis was reported in 29 cases, all of whom were Indian, making the attack rate for that group 53.9 per 100,000 population. Five of the 7 reported cases of tetanus died. A total of 392 cases of German measles was reported, with about 80.0 per cent of those with age specified being under ten years of age.


Other acute communicable disease cases reported were: Mumps 1804, typhemia 35, hookworm 17, Rocky Mountain spotted fever 6, leptospirosis 1, ringworm of the scalp 38, and trichinosis 2. No cases of typhus fever were reported.

RHEUMATIC FEVER

Rheumatic fever cases reported in 1954 were fewer in number than for any year since the disease was made reportable in 1949 in Oklahoma. The number was 52, making a rate of 2.3 for 100,000 population. The attack rate from this disease seems to be on the downward trend but since the time for which data on reported cases are available is so short, the trend line has been compared on Chart 3 on page 8 with the decreasing death rates for the period 1940 to 1954 from acute rheumatic fever and chronic rheumatic heart disease.

The attack rate of this disease among the Indian population was quite high during the year, 20.5 as compared to 1.9 for the white population and 2.1 for the Negro. The six-year average rates, 1949-1954, show somewhat the same relationship, but, of course, big differences due to chance variations from year to year are reduced: 10.2 for the Indian population, 2.9 for the white and 3.2 for the Negro.

Chart 3
Attack Rates from Rheumatic Fever Compared to Death Rates
from Acute Rheumatic Fever and Chronic Rheumatic Heart Disease
Oklahoma, 1940-1954



The rheumatic fever cases reported in 1954 were mostly in children and young adults. About 79 per cent of the cases were under 25 years of age - 58 per cent under 15 years of age. The highest number of cases for any five-year age group was 16 aged 10-14.

TUBERCULOSIS


The tuberculosis attack rate dropped again in 1954 to 73.4 from 78.4 in the previous year, but was still higher than the low rate of 70.3 in 1952. The total number of cases reported was 1,642; 1,601 were specified as respiratory and 41 as other forms. Table 4 on page 9 shows the respiratory cases broken down by stage and activity and the other forms by site. The race-specific attack rate for respiratory tuberculosis was highest among the Indian population, 360.8 per 100,000 population, as compared to 77.7 for the Negro and 57.1 for the white.

Of the 1,526 cases of respiratory tuberculosis with activity specified that were reported for the first time, 44.5 per cent were arrested or of questionable activity. The 55.5 per cent that were specified as active were distributed as follows: 8.7 per cent minimal, 11.6 per cent moderately advanced, 14.7 per cent far advanced, and 17.5 per cent unspecified as to stage. Chart 4 shows a comparison of the three-year average rates for active reported cases specified as minimal, moderately advanced, or far advanced, by age.

Table 4
Reported Cases of Tuberculosis, by Type, Stage
and Activity, by Race, Oklahoma, 1954

Type, Stage and Activity	Total	Race			
		White	Negro	Indian	Unknown
Tuberculosis of respiratory system:					
Minimal, active	133	90	7	20	16
Moderately advanced, active	223	148	18	34	23
Far advanced, active	224	147	23	40	14
Active, unspecified stage	267	186	21	36	24
Arrested (including inactive)	547	446	31	35	35
Activity questionable	132	100	5	12	15
Activity unspecified	75	47	8	17	3
Tuberculosis of other sites:					
Meninges and central nervous system	9	2	-	7	-
Intestines, peritoneum, mesentery	3	3	-	-	-
Vertebral column	1	-	-	1	-
Other bones and joints	10	3	-	5	-
Skin and subcutaneous tissue	6	2	-	2	-
Lymphatic system	1	-	-	-	-
Genito-urinary system	1	-	-	-	-
Adrenal glands	1	-	-	-	-
Other organs	1	-	-	-	-
Disseminated (miliary)	11	5	2	3	1

Chart 4
Reported Case Rates of Active Tuberculosis with Stage Specified,
by Stage and Age, Oklahoma, 1952-1954



VENEREAL DISEASES

The 1954 figure for syphilis cases, reported for the first time, again showed a drop, from 1,989 in 1953 to 1,638. This number was about one-sixth of the number reported during the peak year of 1943, when 9,511 cases were reported. The case rates for the Negro and Indian populations, 369.7 and 256.5, respectively, were high as compared to the white rate of 12.2. About 66 per cent of the reported cases with age specified were 35 years of age and over and 31 per cent were between the ages 15 and 34. Slightly over half of the cases, 51.2 per cent, were in females. Almost three-fourths of the new cases, 73.8 per cent, were diagnosed as being in late or late latent stages of the disease. Table 5 shows the cases by stage for each sex.

Table 5
Reported Cases of Venereal Diseases, by Sex
Oklahoma, 1954

Disease and Stage	Total	Male	Female	Unknown
Total venereal diseases	6,652	3,805	2,804	13
Gonorrhea	4,973	2,972	1,966	35
Syphilis, all stages	1,638	796	834	8
Primary and secondary	48	28	20	-
Early latent	252	87	163	-
Late and late latent	1,209	637	567	5
Congenital	125	41	83	1
Not stated	4	3	1	-
Ophthalmia neonatorum	-	-	-	-
Other venereal diseases	41	37	4	-
Chancroid	38	34	4	-
Granuloma inguinale	1	1	-	-
Lymphogranuloma	2	2	-	-

The number of gonorrhea cases showed an increase over the number for 1953 - 4,973 for 1954 and 4,553 for 1953. Although the number reported annually has dropped from the numbers reported during the years when intensive case-finding programs were in full swing, the drop in gonorrhea cases has not been nearly as great as for syphilis. Almost half as many cases were reported in 1954 as in the year of highest incidence, 1946, when 11,050 cases were reported. The Negro rate for 1954 was 2,010.3 per 100,000 population, as compared to 76.5 for the white and 662.1 for the Indian. The gonorrhea cases were in general much younger than were the syphilis cases: 89 per cent were between the ages 15 and 34 and only 9 per cent were 35 or over. About 60 per cent of the cases were male. Four of the gonorrhea cases were specified as gonococcal ophthalmia and were in infants over four weeks of age but no cases of ophthalmia neonatorum were reported during the year.

Other venereal disease cases numbered 41: chancroid 38, granuloma inguinale 1, and lymphogranuloma 2.

MALIGNANT NEOPLASMS

The reporting of malignant neoplasms is known to be far from complete as evidenced by the number of deaths assigned to this cause which had not been reported as cases during the eight-year period in which cancer has been reportable. Study of the cases that are reported, however, may give some insight into the problems of control of this disease. A total of 1,212 cases were reported during 1954 - at the rate of 53.5 per 100,000 population among the white group, 56.4 among the Negroes, and 35.3 among Indians. Table V in the appendix shows the cases for each sex and each racial group by primary site of the malignant neoplasms.

For the general population, the primary site of reported cases of cancer most frequently specified was the skin, 312 of the cases reported during 1954. The most common primary site for cancer deaths, however, is "digestive organs and peritoneum". Comparison is made in Table 6 of the most frequently reported primary sites for reported cases and for deaths based on three-year average numbers.

Table 6
Most Frequently Reported Primary Site of Malignant Neoplasms
for Cases and Deaths, Oklahoma, 1951-1954

Reported Cases	Average Number 1952-1954	Deaths	Average Number 1951-1953
Skin	349	Digestive organs and peritoneum	1,038
Uterus	174	Respiratory system	292
Digestive organs and peritoneum	173	Uterus	256
Respiratory system	106	Breast	220
Breast	106	Male genital organs	209
Buccal cavity and pharynx	94		

About 54 per cent of the cases reported for the first time during the year 1954 were male. The most frequently reported primary site for the male cases was skin; second and third most common sites were "digestive organs and peritoneum" and "buccal cavity and pharynx". For the female cases, the uterus was the most common primary site, with skin and breast in second and third positions as leading sites.

Over two-thirds, 69.3 per cent, of the reported cases of cancer were in persons 55 years of age and over and for most sites, higher proportions of the reported cases were in persons 55 years of age and over. For cancer of the breast, however, 51.6 per cent of the cases were between ages 25 and 54, and for cancer of the uterus, 53.3 per cent were between the ages 25 and 54. Few cases, only 15, or 1.3 per cent of the total, were under 25 years of age.

At the time of report, metastasis had occurred in 54.2 per cent of the total 476 cases in 1954 for which a report was made as to whether or not metastasis had occurred. This information was not specified, however, on 736, 60.7 per cent of the total number of reported cases. Using the statistics compiled for the three-year period 1952-1954 for the total reports that indicated the metastatic status, 37.1 per cent specified that metastasis had occurred. Since a case registry is not maintained on reported cases, it is not known in how many cases metastasis may have occurred at some time subsequent to the date of the first report of each case. Data for the three-year period based on the status at the time the case was first reported to the State Department of Health, indicated that metastasis had occurred most frequently in cases with primary cancer of the digestive organs and peritoneum, 70.5 per cent of the cases, and of the breast, 60.6 per cent. Approximately 66.7 per cent of the cases with primary cancer of the female genital organs (excluding the uterus) were specified as having metastasis, and 55.4 per cent of cases with primary cancer of respiratory organs were specified as having metastasis; a high percentage, 60.0, of cases with primary lymphosarcoma or reticulosarcoma was specified as having metastasis, but this was based on a very small number of reported cases, 5 for the three-year period. The most frequently reported secondary sites were lymphosarcoma and reticulosarcoma, 31.5 per cent of the cases with metastasis of specified site, digestive organs and peritoneum in 16.7 per cent of the cases, bone in 8.2, and respiratory system in 7.9 per cent of the cases.

About 77.4 per cent, 938, of the 1954 case reports indicated whether or not a biopsy had been performed. Of these, 91.3 per cent specified that there had been a biopsy.

Cancer Cases Reported by Death Certificate Only

During the year 1954, a total of 3,047 deaths were attributed to malignant neoplasms; 85.4 per cent of these deaths, 2,601 in all, had not been previously reported as cases to the State Department of Health during the eight-year period that cancer has been reportable. The number of cases reported only by death certificate was more than twice the number that was reported through regular channels, as indicated by the comparative figures in Table 7 on page 13. The cases reported by death certificate only, however, have not been added to the cases reported through regular channels for total morbidity since the dates of onset and the first diagnoses of these cases are not known. The statistical data regarding these cases reported by death certificate may be used to modify whatever observations were made regarding reported cases.

The most frequently reported primary site for the cases reported only after death by means of the death certificate was "digestive organs and the peritoneum", 33.4 per cent of the total cases reported by this means.

Of the total 2,601 cases reported only by the death certificate, 2,410 were white, 143 were Negro, and 48 were Indian. About 51.7 per cent, 1,346 were male. Approximately 79.3 per cent of these cases were 55 years of age and over as compared to the 69.3 per cent of the cases reported through regular channels who were 55 years of age and over.

Table 7
Cases of Cancer Reported Through Regular Channels and by Death Certificate Only, Number and Per Cent, by Primary Site of Lesion Oklahoma, 1954

Primary Site	Cases Reported Through Regular Channels		Cases Reported by Death Certificate Only	
	Number	Per Cent	Number	Per Cent
Total	1,212	99.7	2,601	100.2
Buccal cavity and pharynx	119	9.8	46	1.8
Digestive organs and peritoneum	159	13.1	868	33.4
Respiratory system	102	8.4	93	3.6
Breast	96	7.9	176	6.8
Uterus	175	14.4	161	6.2
Other female genital organs	34	2.8	67	2.6
Male genital organs	47	3.9	191	7.3
Urinary organs	49	4.0	124	4.8
Skin, except of breast, genital organs, or anus	312	25.7	54	2.1
Brain and other parts of central nervous system	3	0.2	64	2.5
Bone	15	1.2	30	1.2
Lymphosarcoma and reticulosarcoma	10	0.8	46	1.8
Hodgkin's disease	6	0.5	27	1.0
Leukemia and leukemia	12	1.0	152	5.8
Other lymphatic and hematopoietic tissues	4	0.3	32	1.2
Other and unspecified sites	69	5.7	470	18.1

TABLE 1. REPORTED CASES OF SELECTED COMMUNICABLE DISEASES, NUMBER AND RATE, (NUMBER PER 100,000 ESTIMATED POPULATION), OKLAHOMA, 1915-1951

Disease	1915		1916		1917		1918		1919	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Anthrax in man	2	0.2	-	-	1	0.0	86	3.8	114	6.1
Breastlouse	37	1.6	36	1.6	89	3.9	1,117	62.7	1,751	78.1
Chickpox	682	29.9	632	27.8	966	42.7	1,151	62.7	1,321	5.9
Diphtheria	300	13.1	222	9.8	209	9.2	165	7.3	132	5.9
Dysentery	211	9.2	80	3.5	119	6.6	163	7.2	273	12.2
Gonorrhea	8	0.4	24	1.1	13	0.6	8	0.4	12	0.7
Hepatitis, infectious	9,216	403.8	11,050	486.4	9,335	412.7	7,082	314.6	5,987	267.1
Malaria, acquired in U. S.	1,101	46.2	308	13.6	536	23.7	1,021	17.8	86	3.8
Malaria, acquired outside U. S.	811	36.8	4,307	191.1	71	3.1	1,633	72.5	7,538	336.4
Measles	81	3.7	77	3.4	67	3.0	65	2.9	56	2.5
Meningococcal infections	179	7.9	152	6.8	660	29.2	887	39.4	2,764	123.3
Paratyphoid fever	8	0.4	131	5.9	43	1.9	5	0.2	1,352	59.0
Paratyphoid fever, acute	200	8.8	2	0.1	-	-	369	16.4	-	-
Rheumatic fever	-	-	-	-	-	-	-	-	-	-
Rocky Mountain spotted fever	25	1.1	30	1.3	36	1.6	30	1.3	104	4.6
Scarlet fever	1,253	54.9	1,306	58.0	1,353	59.2	1,167	51.3	1,462	64.2
Septic sore throat	124	5.5	180	7.9	198	8.8	176	7.8	308	13.5
Smallpox	13	0.6	16	0.7	4	0.2	1	0.0	2	0.1
Syphilis	5,970	261.9	7,993	347.8	7,117	317.3	5,727	254.4	3,697	163.2
Tuberculosis, all forms	2,678	118.4	2,415	107.7	2,117	94.7	2,346	104.3	2,402	107.2
Typhoid fever	92	4.0	54	2.4	96	4.2	74	3.3	74	3.3
Whooping cough	713	31.2	479	21.1	1,095	48.6	1,084	48.1	228	10.2

Symbols Used in Tables

- Number or rate is zero
- ... Item not applicable
- 0.0 Rate is more than 0 but less than 0.05
- Data not available

Disease	1920		1921		1922		1923		1924	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Anthrax in man	-	-	-	-	62	2.8	1	0.0	10	0.4
Chickpox	2,512	112.5	1,910	85.5	1,211	52.0	1,358	59.2	1,462	64.2
Diphtheria	131	5.9	103	4.6	74	3.3	69	3.1	42	1.9
Dysentery	152	6.8	344	15.4	189	8.5	225	10.1	207	9.2
Gonorrhea	12	0.5	22	1.0	25	1.1	20	0.9	22	1.0
Hepatitis, infectious	5,309	237.7	5,573	249.4	4,823	215.7	4,552	203.5	4,524	202.1
Hepatitis, infections	-	-	73	3.3	70	3.1	115	5.1	317	14.5
Malaria, acquired in U. S.	92	4.1	14	0.6	12	0.5	11	0.5	19	0.8
Malaria, acquired outside U. S.	2	0.1	372	16.8	11	0.5	11	0.5	12	0.5
Measles	448	20.0	8,000	358.0	76	3.4	3,798	169.8	3,403	152.3
Meningococcal infections	56	2.5	72	3.2	56	2.5	54	2.4	58	2.6
Mumps	2,937	131.5	1,659	74.2	865	38.3	898	40.2	1,044	46.6
Paratyphoid fever	14	0.6	67	3.0	39	1.7	68	3.0	59	2.6
Polioomyelitis, acute	533	23.9	8	0.4	2	0.1	536	24.0	559	25.0
Rabies in man	-	-	-	-	-	-	-	-	-	-
Rheumatic fever	73	3.3	82	3.7	58	2.6	57	2.5	52	2.3
Rocky Mountain spotted fever	10	0.4	7	0.3	1	0.0	10	0.4	10	0.4
Scarlet fever	532	23.8	196	8.8	382	17.1	684	30.6	736	32.9
Septic sore throat	387	17.3	532	23.8	277	12.4	262	11.7	253	11.3
Smallpox	5	0.2	2,906	130.0	2,351	105.2	1,989	88.9	1,658	73.2
Syphilis	3,169	141.9	2,090	93.8	1,762	78.4	1,754	78.4	1,682	73.4
Tuberculosis, all forms	2,090	90.9	2,172	96.8	38	1.7	46	2.1	35	1.6
Typhoid fever	84	3.8	53	2.4	61	2.7	46	2.1	43	1.9
Typhus fever	1	0.0	-	-	372	16.6	256	11.4	219	9.8
Whooping cough	933	41.8	1,115	49.9	-	-	-	-	-	-

TABLE IV. REPORTED CIVILIAN CASES OF COMMUNICABLE DISEASES, NUMBER AND RATE (NUMBER PER 100,000 ESTIMATED POPULATION) AND NUMBER BY RACE, BY URBAN AND RURAL RESIDENCE, OKLAHOMA, 1951.

	Estimated population, 7-1-54	Total	Urban				Total				Rural			
			Total		Race		Total		Race		Total		Race	
			No.	Ratio	No.	No.	No.	Ratio	No.	No.	No.	No.	No.	No.
Attack in man		10	-	0.7	-	13	11	125	31	3.0	52	-	36	-
Breast-labs		1,382	953	69.7	792	13	125	121	21	12.5	49	5	48	3
Cholera		19	23	1.2	18	7	7	12	12	1.5	38	1	37	1
Dysentery, amebic		24	0	0.6	5	2	-	17	17	1.6	16	1	15	1
Dysentery, bacillary		155	63	6.9	26	3	54	73	7.3	7.1	29	2	27	2
Dysentery, unspecified		13	13	1.2	10	2	1	12	12	1.2	12	1	11	1
Enteric fever		17	11	1.4	10	2	1	12	12	1.2	12	1	11	1
German measles		360	293	24.3	198	2	92	67	6.5	6.1	61	4	57	4
Gonorrhea		4,193	3,107	288.2	1,075	2,609	532	733	69.9	263	30.5	102	102	102
Impetigo, intertrigo		334	175	14.5	104	1	4	15	15.4	14	11	11	11	11
Measles		15	15	1.2	14	5	4	15	15.4	14	11	11	11	11
Malaria, acquired outside U. S.		6	1	0.2	1	15	32	1,032	618	62.7	16	1	15	1
Malaria, acquired in U. S.		3,346	2,666	224.1	1,633	5	13	256	681	60.3	147	72	54	54
Non-specific infections		1,736	1,112	92.4	836	7	256	681	60.3	147	72	54	54	54
Opportunistic neontoma		58	34	3.4	34	8	1	2	21	2.0	20	2	18	2
Paratyphoid fever		58	34	3.4	34	8	1	2	21	2.0	20	2	18	2
Scarlet fever, acute		52	27	2.2	21	3	-	25	24	2.4	17	-	17	-
Rheumatic fever		52	27	2.2	21	3	-	25	24	2.4	17	-	17	-
Rocky Mountain spotted fever		73	56	4.6	25	3	10	12	10.4	13	1	-	1	-
Sarcoma		534	309	25.7	214	8	6	51	25	2.5	207	9	198	9
Serum sore throat		1,610	1,202	100.4	651	69	58	101	30.8	203	11.6	70	12	58
Syphilis		69	5	0.4	4	1	-	21	21	2.1	20	-	19	-
Tuberculosis		1,601	892	74.1	656	76	83	77	709	68.6	508	37	471	37
Tuberculosis, respiratory		14	28	2.3	21	3	1	3	21	2.1	2	1	1	1
Tuberculosis, other forms		63	23	1.9	21	1	1	10	10	1.0	59	1	58	1
Typhoid fever		30	21	1.7	8	11	1	2	2	0.2	5	3	2	3
Typhus fever		21	21	1.7	8	11	1	2	2	0.2	5	3	2	3
Unspecified fever		150	150	12.6	110	26	7	67	6.5	48	48	1	47	1
Whooping cough		219	150	12.6	110	26	7	67	6.5	48	48	1	47	1

TABLE V. REPORTED CASES OF HALLMARK NEOPLASMS, BY PRIMATE SITE OF LESION, RACE AND SEX, OKLAHOMA, 1954

Primary Site	Total	Sex		Race			
		Male	Female	White	Negro	Indian	Unknown
Total, all sites	1,312	652	560	1,091	82	19	20
Breast cavity and pharynx	112	97	22	112	3	2	2
Larynx and trachea	118	112	6	112	10	3	3
Esophagus and peritoneum	112	97	15	112	8	—	—
Respiratory system	102	86	16	91	10	2	3
Breast	96	1	95	61	3	2	—
Uterus	175	—	175	132	32	6	5
Other female genital organs	115	—	31	30	—	—	—
Male genital organs	147	147	—	147	7	1	1
Urinary organs	147	37	12	—	—	—	—
Skin, except of breast, genital organs, or anus	312	210	102	304	3	3	2
Bone and central nervous system	112	102	—	112	—	—	—
Bone	112	10	5	12	2	—	—
Lymphatic and reticuloendothelium	10	8	2	10	—	—	—
Hodgkin's disease	6	—	—	—	—	—	—
Leukemia and aplasia	12	9	3	11	1	—	—
Other lymphatic and hematopoietic	—	—	—	—	—	—	—
Stomach	49	2	26	1	2	1	1
Other and unspecified sites	4	1	—	4	—	—	—

TABLE VI. REPORTED CASES OF SELECTED COMMUNICABLE DISEASES BY SEX AND RACE
OKLAHOMA, 1954

Disease	Total			White			Negro			Indian			Unknown		
	Male	Female	Unknown	Male	Female	Unknown	Male	Female	Unknown	Male	Female	Unknown	Male	Female	Unknown
Anthrax in man	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Brucellosis	32	8	-	30	8	-	-	-	-	2	-	-	-	-	-
Chickenpox	648	591	176	600	553	10	15	8	-	28	28	-	5	2	166
Diphtheria	24	18	-	20	16	-	3	2	-	1	-	-	-	-	-
Dysentery	100	105	2	41	48	-	7	8	-	50	48	-	2	1	2
Encephalitis, infectious	13	11	-	10	10	-	1	1	-	-	-	-	2	-	-
German measles	171	129	92	165	125	-	2	1	-	3	2	-	1	1	92
Gonorrhea	2,972	1,966	35	910	690	-	1,895	1,030	-	134	222	-	33	24	35
Hepatitis, infectious	168	173	6	157	158	-	6	3	-	5	11	-	-	1	6
Hookworm	10	7	-	3	2	-	1	-	-	6	5	-	-	-	-
Malaria, acquired in U. S.	10	9	-	10	8	-	-	-	-	1	-	-	-	-	-
Malaria, acquired outside U. S.	6	1	-	6	1	-	-	-	-	-	-	-	-	-	-
Measles	1,164	1,058	1,181	1,047	982	131	32	12	-	28	28	5	57	36	1,045
Meningococcal infections	27	31	-	24	27	-	2	4	-	1	-	-	-	-	-
Mumps	808	600	396	725	551	58	26	6	64	41	26	-	16	17	274
Ophthalmia neonatorum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Paratyphoid fever	23	15	-	19	15	-	-	-	-	2	-	-	2	-	-
Polycycolitis, acute	323	235	1	311	223	-	3	7	-	7	4	-	2	1	1
Rheumatic fever	30	22	-	20	18	-	3	-	-	7	4	-	-	-	-
Rocky Mountain spotted fever	2	4	-	1	3	-	-	-	-	-	1	-	1	-	-
Scarlet fever	371	338	27	366	323	10	3	5	-	1	6	-	1	4	17
Septic sore throat	280	208	78	262	195	20	13	10	-	5	3	-	-	-	58
Smallpox	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Syphilis	796	834	8	446	415	-	264	302	1	52	87	-	34	30	7
Tetanus	5	2	-	1	-	-	4	2	-	-	-	-	-	-	-
Trachoma	13	16	-	-	-	-	-	-	-	13	16	-	-	-	-
Tuberculosis, respiratory	991	599	11	733	429	2	70	43	-	111	82	1	77	45	8
Tuberculosis, other forms	25	14	2	10	5	-	3	1	-	10	8	-	2	-	2
Tularemia	24	11	-	20	10	-	1	-	-	2	1	-	1	-	-
Typhoid fever	41	22	-	31	19	-	1	-	-	9	2	-	-	1	-
Typhus fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Venereal diseases, other	37	4	-	18	1	-	16	3	-	2	-	-	1	-	-
Vincent's angina	10	12	-	5	7	-	-	-	-	2	-	-	5	3	-
Whooping cough	93	109	17	73	91	10	14	13	-	6	5	-	-	-	7

TABLE VII. REPORTED CASES OF SELECTED COMMUNICABLE DISEASES BY AGE
OKLAHOMA, 1954

Disease	All Ages	Age in Years																	Unknown
		Under 1 year	1	2	3	4	5-9	10-14	15-19	20-24	25-29	30-34	35-44	45-54	55-64	65-74	75-84	85 and Over	
Anthrax in man	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Brucellosis	40	-	-	-	-	-	-	-	2	1	5	4	10	6	2	3	1	-	-
Chickenpox	1,415	60	74	79	114	122	632	53	16	27	17	5	8	1	2	-	-	-	205
Diphtheria	42	-	6	3	3	1	7	6	5	1	4	2	3	-	-	-	-	-	1
Dysentery	207	41	24	22	11	3	22	15	9	7	6	8	6	9	5	4	3	1	11
Encephalitis, infectious	24	1	-	3	-	-	2	3	-	1	-	-	3	3	5	-	1	-	2
German measles	392	61	42	14	17	23	79	8	27	20	4	-	-	-	-	-	-	-	97
Gonorrhea	4,973	13	3	3	3	2	17	68	1,304	1,862	787	371	306	95	22	10	2	-	105
Hepatitis, infectious	347	-	-	2	8	12	106	69	26	20	22	18	4	5	4	-	-	1	29
Hookworm	17	-	-	1	1	1	4	5	4	-	-	-	1	-	-	-	-	-	-
Malaria, acquired in U. S.	19	-	-	-	-	-	-	-	1	2	3	4	3	3	-	2	1	-	-
Malaria, acquired outside U. S.	7	-	-	-	-	-	-	-	-	3	3	1	-	-	-	-	-	-	-
Measles	3,403	121	185	171	192	221	909	77	49	25	11	2	11	1	-	-	1	-	1,427
Meningococcal infections	58	10	5	6	4	5	12	7	-	-	-	2	4	1	1	1	-	-	-
Mumps	1,804	6	24	42	55	80	503	149	137	79	62	71	76	23	6	6	3	-	482
Ophthalmia neonatorum	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Paratyphoid fever	38	1	1	-	-	-	2	1	1	3	3	5	13	2	1	1	2	-	2
Polycycolitis, acute	559	19	30	41	35	33	140	94	42	40	35	19	19	3	4	-	-	-	5
Rheumatic fever	52	-	-	1	1	3	9	16	9	2	2	3	2	3	-	-	1	-	-
Rocky Mountain spotted fever	6	-	-	-	-	1	1	1	-	-	-	-	1	-	-	1	-	-	1
Scarlet fever	736	9	28	33	60	63	380	81	17	5	1	3	1	2	1	-	-	-	52
Septic sore throat	566	3	16	18	31	34	94	47	46	54	33	37	32	20	4	2	1	1	93
Smallpox	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Syphilis	1,638	5	-	-	1	-	2	16	90	120	122	118	333	329	225	87	21	1	168
Tetanus	7	2	-	2	-	-	-	-	-	-	-	-	1	-	-	-	1	-	1
Trachoma	29	-	-	-	-	-	6	13	3	-	-	-	-	1	2	4	-	-	-
Tuberculosis, respiratory	1,601	1	3	5	6	5	17	26	58	101	98	117	202	284	294	193	90	12	89
Tuberculosis, other forms	41	2	3	2	-	1	1	1	2	5	1	3	4	4	6	1	2	1	2
Tularemia	35	-	-	-	1	-	2	4	2	1	3	1	6	7	2	2	1	-	3
Typhoid fever	63	-	1	2	-	2	8	11	9	3	3	3	6	7	-	3	1	-	4
Typhus fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Veneral diseases, other	41	-	-	-	-	-	-	1	7	16	8	1	3	2	-	2	-	-	1
Vincent's angina	22	-	1	-	-	-	1	4	5	2	-	-	1	-	-	-	-	-	8
Whooping cough	219	40	23	19	22	14	57	21	3	1	-	-	-	1	-	-	-	-	18

TABLE VIII. REPORTED CASES OF COMMUNICABLE DISEASES BY COUNTY OF RESIDENCE
OKLAHOMA CITY, TULSA CITY AND MIDLAND, 1954

Disease	State	Adair	Alfalfa	Atoka	Beaver	Beckham	Blaine	Butte	Caddo	Canadian
Anthrax in man	10	-	-	-	-	-	-	-	-	-
Brucellosis	1,125	12	7	5	1	8	12	1	1	1
Diphtheria	24	-	-	-	-	-	-	-	-	-
Dysentery, amebic	156	-	-	-	-	-	-	-	-	-
Dysentery, bacillary	24	-	-	-	-	-	-	-	-	-
Dysentery, unspecified	292	23	1	1	1	-	59	27	52	15
German measles	4,372	17	-	-	-	-	-	-	-	-
Gonorrhea	373	22	1	11	-	-	-	-	-	-
Hepatitis, infectious	17	-	-	-	-	-	-	-	-	-
Histoplasmosis	19	-	-	-	-	-	-	-	-	-
Measles, acquired in U. S.	3,403	10	-	-	-	-	-	-	-	-
Measles, acquired outside U. S.	1,604	21	-	-	-	-	-	-	-	-
Meningococcal infections	38	-	-	-	-	-	-	-	-	-
Opthalmia neonatorum	559	2	-	-	-	-	-	-	-	-
Paratyphoid fever	54	-	-	-	-	-	-	-	-	-
Poliovirus, acute	52	-	-	-	-	-	-	-	-	-
Rabies in animals	6	-	-	-	-	-	-	-	-	-
Rabies in man	566	-	-	-	-	-	-	-	-	-
Rheumatic fever	1,659	13	-	-	-	-	-	-	-	-
Scarlet fever	29	-	-	-	-	-	-	-	-	-
Scarlet sore throat	1,401	23	-	-	-	-	-	-	-	-
Syphilis	63	-	-	-	-	-	-	-	-	-
Tetanus	15	-	-	-	-	-	-	-	-	-
Tuberculosis, respiratory	1,401	23	-	-	-	-	-	-	-	-
Tuberculosis, other forms	15	-	-	-	-	-	-	-	-	-
Typhoid fever	63	-	-	-	-	-	-	-	-	-
Typhus fever	12	-	-	-	-	-	-	-	-	-
Unusual diseases, other	219	-	-	-	-	-	-	-	-	-
Whooping cough	-	-	-	-	-	-	-	-	-	-

TABLE VIII. REPORTED CASES OF COMMUNICABLE DISEASES BY COUNTY OF RESIDENCE
OKLAHOMA CITY, TULSA CITY AND MIDLAND, 1954
(Continued)

Disease	Osage	Delaware	DeWey	Ellis	Garfield	Guthrie	Harper	Hotchkiss	Hughes	Jackson	Jeff-	Johnston	Key	King-	Knox	Latimer
Anthrax in man	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Brucellosis	33	2	4	4	40	17	4	1	2	2	3	5	19	1	29	3
Diphtheria	1	1	-	-	1	-	-	-	-	-	-	-	-	-	-	-
Dysentery, amebic	11	-	1	1	21	4	-	-	-	-	-	-	-	-	-	-
Dysentery, bacillary	1	-	1	1	1	1	-	-	-	-	-	-	-	-	-	-
Dysentery, unspecified	5	1	1	1	16	11	-	-	-	-	-	-	-	-	-	-
German measles	60	5	4	2	29	11	-	-	-	-	-	-	-	-	-	-
Gonorrhea	2	-	-	-	2	0	-	-	-	-	-	-	-	-	-	-
Hepatitis, infectious	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Histoplasmosis	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Measles, acquired in U. S.	153	-	-	-	271	2	-	-	-	-	-	-	-	-	-	-
Measles, acquired outside U. S.	12	-	-	-	119	2	-	-	-	-	-	-	-	-	-	-
Meningococcal infections	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Opthalmia neonatorum	19	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Paratyphoid fever	14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Poliovirus, acute	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rabies in animals	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rabies in man	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rheumatic fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scarlet fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scarlet sore throat	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Syphilis	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tetanus	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tuberculosis, respiratory	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tuberculosis, other	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Typhoid fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Typhus fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unusual diseases, other	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Whooping cough	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Disease	Osage	Delaware	DeWey	Ellis	Garfield	Guthrie	Harper	Hotchkiss	Hughes	Jackson	Jeff-	Johnston	Key	King-	Knox	Latimer
Anthrax in man	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Brucellosis	7	13	8	20	43	3	4	1	1	2	3	5	19	1	29	3
Diphtheria	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Dysentery, amebic	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Dysentery, bacillary	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Dysentery, unspecified	3	2	62	10	20	24	1	1	1	1	1	1	1	1	1	1
German measles	51	2	62	11	24	1	1	1	1	1	1	1	1	1	1	1
Gonorrhea	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hepatitis, infectious	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Histoplasmosis	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Measles, acquired in U. S.	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Measles, acquired outside U. S.	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Meningococcal infections	21	6	26	13	35	14	22	11	13	34	7	28	72	11	15	2
Opthalmia neonatorum	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Paratyphoid fever	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Poliovirus, acute	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rabies in animals	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rabies in man	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rheumatic fever	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scarlet fever	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scarlet sore throat	11	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Syphilis	29	10	24	1	27	4	35	4	4	10	4	4	4	4	4	4
Tetanus	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tuberculosis, respiratory	36	19	24	1	21	2	27	7	11	32	10	20	21	2	12	2
Tuberculosis, other forms	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Typhoid fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Typhus fever	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unusual diseases, other	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Whooping cough	4	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-

TABLE VIII. REPORTED CASES OF COMMUNICABLE DISEASES BY COUNTY OF RESIDENCE
OKLAHOMA CITY, TULSA CITY AND MILWAUKEE, 1954
(Continued)

Disease	LaFlore	Lincoln	Logan	Lone	Medan	McCurrah	McNabb	Major	Marshall	Wyer
Anthrax in man	1	1	-	-	-	-	-	-	-	-
Bubonic plague	17	11	13	-	5	3	-	13	4	9
Dysentery, amebic	-	-	-	-	-	12	-	2	-	-
Dysentery, bacillary	-	-	-	-	-	57	-	-	-	-
Epidemic typhus	-	-	6	6	-	61	-	-	-	1
Enteric fever	-	-	1	1	-	6	-	-	-	-
German measles	22	11	131	1	1	6	37	1	13	6
Gonorrhea	2	-	1	-	-	-	-	1	-	1
Histoplasmosis, infectious	-	-	-	-	-	-	-	-	-	-
Measles, acquired in U. S.	-	-	-	-	-	-	-	-	-	-
Measles, acquired outside U. S.	-	-	15	-	22	150	2	32	3	39
Meningococcal infections	4	4	114	-	10	98	2	7	1	31
Scarlet fever	-	-	-	-	-	-	-	-	-	-
Smallpox	15	5	8	-	3	20	14	-	4	4
Typhoid fever	-	-	-	-	-	-	-	-	-	-
Typhus	74	5	1	1	3	32	14	-	4	1
Typhus, abdominal	36	17	22	6	4	32	15	-	2	24
Typhus, respiratory	1	-	-	-	-	-	-	-	-	-
Typhus, enteric	7	-	-	-	-	-	-	-	-	-
Typhus, paratyphoid	4	-	-	-	-	-	-	-	-	-
Typhus, other forms	14	-	-	-	-	-	-	-	-	-
Typhus fever	7	-	-	-	-	-	-	-	-	-
Varicella	1	-	-	-	-	-	-	-	-	-
Viral diseases, other	-	-	-	-	-	-	-	-	-	-
Vincennes, angina	-	-	-	-	-	-	-	-	-	-
Whooping cough	-	5	23	-	-	2	1	-	1	5

TABLE VIII. REPORTED CASES OF COMMUNICABLE DISEASES BY COUNTY OF RESIDENCE
OKLAHOMA CITY, TULSA CITY AND HILTIARY, 1954
(Continued)

Disease	Egypto	Pitt- tong	Pontoto	Poke- toke	Park- madam	Roger Hills	Ingore	Semotale	Semogah	Stopham
Anthrax in man	2	2	-	-	1	1	1	-	1	-
Breast cancer	33	65	35	4	-	1	-	2	1	6
Diphtheria	1	1	1	-	-	1	-	-	-	-
Dysentery, amebic	1	1	1	-	1	6	-	-	-	-
Dysentery, bacillary	-	-	-	-	-	-	-	-	-	-
Epidemic typhus	-	-	-	-	-	-	-	-	-	-
Exanthematous diseases	-	-	-	-	-	-	-	-	-	-
German measles	6	42	18	59	-	-	-	17	17	34
Hepatitis	38	52	44	4	1	-	14	10	-	4
Hookworm	1	24	5	1	1	-	-	-	-	-
Intestinal diseases	-	-	-	-	-	-	-	-	-	-
Measles, acquired in U. S.	-	-	-	-	-	-	-	-	-	-
Malaria, acquired outside U.S.	14	-	170	68	-	1	2	-	1	18
Meningeal infections	11	71	-	-	-	1	1	-	-	-
Paratyphoid fever	19	19	50	89	1	1	3	1	1	1
Paratyphoid fever	7	18	3	14	1	1	1	1	1	1
Polio, poliomyelitis, acute	-	-	-	-	-	2	2	-	-	12
Rabies in animals	-	-	-	-	-	-	-	-	-	-
Rheumatic fever	-	1	-	-	-	-	-	-	2	-
Rocky Mountain spotted fever	-	-	-	-	-	-	-	-	-	-
Scurvy	10	36	2	8	-	-	1	8	1	8
Schistosomiasis	-	-	-	-	-	1	-	-	-	-
Syphilis	9	5	2	-	-	1	1	1	1	1
Typhoid fever	9	47	17	36	3	9	12	26	9	17
Tuberculosis	22	25	19	32	9	2	2	2	39	16
Tuberculosis, respiratory	-	-	-	-	-	-	-	-	-	-
Typhoid fever	1	1	1	1	1	1	-	2	2	1
Typhus fever	-	1	-	-	-	-	-	-	1	1
Various diseases	1	1	-	1	-	-	-	-	-	-
Whooping cough	1	2	-	1	-	1	-	-	-	-

TABLE VIII. REPORTED CASES OF COMMUNICABLE DISEASES BY COUNTY OF RESIDENCE
OKLAHOMA CITY, TULSA CITY AND MILWAUKEE, 1954
(Continued)

Disease	Hurray	putogase	Hoble	Howata	Ockwison	Ockwison	Oomage	Osage	Oskan	Pamne
Amber in man	-	1	1	3	-	1	1	-	2	1
Brucellosis	18	18	26	3	4	170	4	35	25	17
Cholera	-	-	-	-	-	6	1	-	-	1
Dysentery, amebic	1	-	-	-	-	1	1	-	-	-
Dysentery, bacillary	1	-	-	1	-	12	1	-	1	1
Dysentery, unspecified	6	-	-	-	-	6	-	-	-	-
Enteric fever	-	1	-	-	-	4	1	-	-	-
Enteritis	1	1	-	-	-	h	-	-	-	-
Gonorrhea	24	24	24	5	12	1,423	30	6	1	2
Hepatitis, infectious	35	35	-	1	8	62	209	11	9	2
Marasmus	-	-	-	-	-	-	-	-	-	3
Malaria, acquired in U. S.	-	-	-	-	-	-	-	-	-	2
Malaria, acquired outside U. S.	-	-	-	-	-	-	-	-	-	-
Measles	1	23	63	23	-	164	42	15	90	-
Mumps	-	3	20	-	-	24	15	13	4	9
Ophthalmic neotrom	32	32	10	1	3	285	1	1	1	-
Paratyphoid fever	-	1	-	-	-	h	-	-	-	-
Paratyphoid fever, enteric	-	26	2	1	3	76	4	2	7	8
Rabies in animals	-	-	-	2	-	-	-	-	-	-
Rabies in man	-	3	1	2	-	7	-	1	1	2
Rheumatic fever	-	-	-	-	-	-	-	-	-	-
Rheumatic fever, acute	-	2	-	-	-	2	-	-	-	-
Rheumatic fever, chronic	-	3	1	3	-	7	-	-	-	-
Scarlet fever	-	2	15	-	-	124	17	1	1	1
Septic sore throat	-	-	-	-	-	2	-	1	2	6
Syphilis	-	2	-	-	-	40	129	27	10	12
Tetanus	16	123	5	12	16	380	43	18	17	-
Typhemia	4	4	-	-	-	-	-	-	-	-
Typhoid fever	1	1	20	18	12	207	30	3	85	12
Typhoid fever, enteric	1	1	-	1	2	6	2	2	1	-
Typhoid fever, other forms	2	2	-	-	-	5	-	-	-	-
Typhus fever	-	-	-	-	-	12	-	-	-	-
Typhus fever, unspecified	-	-	-	-	-	12	-	-	-	-
Vaccinia, other	-	-	-	-	-	1	-	-	-	-
Vaccinia, smallpox	-	-	-	-	-	1	-	-	-	-
Vaccinia, other	-	-	-	-	-	30	-	-	-	-
Whooping cough	-	4	-	-	-	1	6	6	1	2

TABLE VIII. REPORTED CASES OF COMMUNICABLE DISEASES BY COUNTY OF RESIDENCE
OKLAHOMA CITY, TULSA CITY AND HILTIARY, 1954
(Continued)

Disease	Texas	Tillman	Tulsa	Wagoner	Maple- lurgon	Wheeler	Hood	Hood- ward	W.H.- kay	Osage City	Tulsa City
Anthrax in man	-	-	2	-	-	-	1	-	-	1	-
Brucellosis	2	3	12	-	20	12	9	40	33	130	109
Diphtheria	-	1	3	-	-	1	1	-	-	109	3
Dysentery, bacillary	-	1	2	-	-	6	-	-	-	10	1
Dysentery, amoebic	-	2	1	-	2	-	-	3	-	1	2
Hepatitis, infectious	-	-	2	-	-	-	-	-	-	1	-
Neurophallitis, infectious	1	-	-	-	-	-	-	-	-	-	-
Ornithion menitis	-	-	-	-	-	-	-	-	-	-	-
Scabies	-	-	1	-	-	-	2	-	-	32	15
Scrofula	-	-	12	-	-	1	2	-	32	15	18
Syphilis	-	-	295	21	13	7	4	2	2	780	264
Hookworm	35	-	24	-	-	-	4	1	13	10	24
Leishmaniasis	-	-	13	-	-	-	-	-	-	-	-
Malaria, acquired in U. S.	2	-	-	-	-	-	-	1	1	1	1
Malaria, acquired outside U. S.	3	1,055	3	1	23	10	5	22	1	1	12
Meningococcal infections	-	-	115	-	7	11	25	57	68	1,040	122
Mumps	6	-	-	-	-	10	12	22	57	67	22
Optic atrophy	-	-	-	-	-	-	-	-	-	-	-
Optic neuritis	-	1	-	-	-	-	-	-	-	-	-
Poliomyelitis, acute	7	1	31	1	20	-	2	13	4	97	99
Rabies in animals	-	1	20	-	1	-	-	-	-	1	1
Rabies in man	-	-	-	-	-	-	-	-	-	-	-
Rocky Mountain spotted fever	5	-	3	-	-	-	-	3	-	3	3
Sand fever	35	277	1	-	1	2	13	3	1	10	29
Septic sore throat	-	20	-	-	3	7	-	21	32	22	10
Scarlet fever	-	-	-	-	-	-	-	-	-	-	-
Scrub typhus	-	-	-	-	-	-	-	-	-	-	-
Syphilis	3	24	206	19	16	2	3	15	28	355	168
Tetanus	-	-	-	-	-	-	-	-	-	-	-
Trenchna	-	-	-	-	-	-	-	-	-	-	-
Tuberculosis, respiratory	-	7	144	-	31	1	2	15	-	155	194
Tuberculosis, other forms	-	-	-	-	-	-	-	-	-	6	-
Typhoid fever	-	-	-	-	1	-	-	-	-	1	1
Dysma fever	-	-	-	-	-	-	-	-	-	1	-
Typhus fever	-	-	-	-	-	-	-	-	-	11	11
Visceral leishmaniasis, other	-	-	-	-	-	-	-	-	-	4	4
Visceral leishmaniasis, other	-	-	-	-	-	-	-	-	-	11	11
Whooping cough	-	28	-	-	1	3	1	-	-	23	23

1. The first part of the document is a list of the names of the persons who were present at the meeting. The names are listed in alphabetical order.

2. The second part of the document is a list of the topics that were discussed at the meeting. The topics are listed in alphabetical order.

3. The third part of the document is a list of the actions that were taken at the meeting. The actions are listed in alphabetical order.