# CLINICAL REMINDERS FOR PRESCRIBING OPIOIDS















**Learn more:** oklahoma.gov/health/overdose

This publication was supported by the Cooperative Agreement Number 5 NOT/CEU02/45-02 funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services. This publication was issued by the Oklahoma State Department of Health (OSDH), an equal opportunity employer and provider. A digital file has been deposited with the Publications Clearinghouse of the Oklahoma Department of Libraries in compliance with section 3-114 of Title 65 of the Oklahoma Statutes and is available for download at documents.ok.gov. I Issued May 2023. Adapted with permission from the Centers for Disease Control and Prevention.

# **Clinical Reminders**



### **Determining When to Initiate or Continue Opioids For Chronic Pain**

- Opioids are not first-line or routine therapy for chronic pain
- Establish and measure goals for pain and function
- Discuss benefits, risks, and availability of nonopioid therapies with patient



# Opioid Selection, Dosage, Duration, Follow-Up and Discontinuation

- Use immediate-release opioids when starting
- Start low and go slow
- When opioids are needed for acute pain, prescribe no more than needed
- Do not prescribe ER/LA opioids for acute pain
- Follow-up and re-evaluate risk of harm; reduce dose or taper and discontinue if needed



# **Assessing Risk and Addressing Harms of Opioid Use**

- Evaluate risk factors for opioid-related harms
- : Check PMP for high dosages and prescriptions from other providers
- Use urine drug testing to identify prescribed substances and undisclosed use
- Avoid concurrent benzodiazepine and opioid prescribing
- Arrange treatment for opioid use disorder if needed